



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

Remark on solution of Problem 299, by J. M. ARNOLD, Crompton, R. I.

The following method shows *why* there is only *one* solution, a fact not clearly discernable in the published solution on page 203, Vol. XV.

Let x , y , $3y-x$ be the sides in increasing order of magnitude. Then the area must be $2y-2x$. This leads to an equation of the first degree in x , from which $x = \frac{1}{2}y + \frac{16y}{3y^2 + 16}$.

Substituting for y , the numbers 1, 2, 3, etc., we find $y=4$, $x=3$. All other integral values of y will give fractional values for x , as the second term soon becomes less than $\frac{1}{2}$, and continues to diminish as y increases.

Hence, 3, 4, 5 are the sides of the only triangle satisfying the problem.

302. Proposed by PROF. R. D. CARMICHAEL, Anniston, Ala.

Prove that the system of equations

$$xu + 5yv = 2,$$

$$xv + yu = 1,$$

have no integral solution except one of the unknowns be zero.

I. Solution by E. B. ESCOTT, Ann Arbor, Mich.

Squaring and adding, and multiplying second equation by 5,

$$(xu - 5yv)^2 + 5(xv + yu)^2 = 9.$$

But, since $(xu - 5yv)^2 + 5(xv + yu)^2 = (xu + 5yv)^2 + 5(xv - yu)^2$, we must have the equation

$$X^2 + 5Y^2 = 9,$$

satisfied for two sets of values (X , Y).

The only solutions of this last equation are

$$X = \pm 2, \pm 3, \quad Y = \pm 1, 0,$$

i. e.,

$$\begin{aligned} xu - 5yv &= 2, & xv + yu &= 1, \\ xu + 5yv &= \pm 3, & xv - yu &= 0. \end{aligned}$$

Whence, $2xu=5$ or -1 , $2xv=1$, evidently impossible.

Therefore, the only possible solutions are

$$xu - 5yv = 2, \quad xu + 5yv = \pm 2.$$

Whence, $2xu = 4$ or 0 , $10yv = 0$ or -4 ; i. e., either x , u , y , or v equals zero.

Solved, similarly, by G. B. M. Zerr and V. M. Spunar.

II. Solution by O. C. CARMICHAEL, Oxford, Ala.

If the square of the first equation be added to five times the square of the second equation, we have

$$x^2u^2 + 5x^2v^2 + 25y^2v^2 + 5y^2u^2 = 9.$$

Therefore, there is no integral solution in x , y , u , v except when one of the unknowns is zero; for, if they were all positive integers, $25y^2v^2$ itself would be greater than 9.

303. Proposed by PROF. R. D. CARMICHAEL, Anniston, Ala.

Evaluate the determinant

$$\begin{vmatrix} D_1 & x_1x_2 & x_1x_3 & \dots & x_1x_n \\ x_1x_2 & D_2 & x_2x_3 & \dots & x_2x_n \\ x_1x_3 & x_2x_3 & D_3 & \dots & x_3x_n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ x_1x_n & x_2x_n & x_3x_n & \dots & D_n \end{vmatrix}$$

Solution by J. W. CLAWSON, Ursinus College, Collegeville, Pa.

$$\begin{aligned} \Delta &= x_1x_2x_3\dots x_n \begin{vmatrix} D_1/x_1 & x_2 & x_3 & \dots & x_n \\ x_1 & D_2/x_2 & x_3 & \dots & x_n \\ x_1 & x_2 & D_3/x_3 & \dots & x_n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ x_1 & x_2 & x_3 & \dots & D_n/x_n \end{vmatrix} \\ &= \prod_1^n x_1 \begin{vmatrix} (D_1 - x_1^2)/x_1 & 0 & 0 & \dots & (x_n^2 - D_n)/x_n \\ 0 & (D_2 - x_2^2)/x_2 & 0 & \dots & (x_n^2 - D_n)/x_n \\ 0 & 0 & (D_3 - x_3^2)/x_3 & \dots & (x_n^2 - D_n)/x_n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ x_1 & x_2 & x_3 & \dots & D_n/x_n \end{vmatrix} \end{aligned}$$

Subtracting the last row from each row,